

LIST OF FIGURES

| Figure | Page |
|---|-------------|
| LITERATURES REVIEW | |
| 1 Appearance of <i>Bordetella pertussis</i> after staining by Gram stain | 6 |
| 2 Colonization of tracheal epithelial cells by <i>Bordetella pertussis</i> | 9 |
| 3 The pathogenesis of pertussis | 10 |
| 4 The NMR structure of the pertussis toxin | 12 |
| 5 The ability of pertussis toxin to interfere the regulation of the eukaryotic adenylate cyclase complex | 14 |
| 6 Schematic representation of an intact Ig together with Fab and Fv fragments and single V and C domains | 27 |
| 7 Schematic representation of the humanization protocols of VL region of antibody | 30 |
| 8 The principle of phage display | 36 |
| 9 The phage display cycle | 37 |
| 10 Phage display formats | 40 |
| 11 Structure of the T7 phage particle | 45 |
| 12 Phage display vectors and the genetic map of T7 | 49 |
| 13 The general stages of a filamentous bacteriophage | 50 |
| MATERIALS AND METHODS | |
| 14 Schematic of production of humanized-PT6-2G6-HuScFv coding sequence | 69 |
| RESULTS | |
| 15 Western blot analysis of MAbPT6-2G6 against electrophoresed-PT | 80 |
| 16 Hemagglutination assay for determining HA100 of PT and hemagglutination inhibition assay for determining the HI activity of MAbPT6-2G6 | 81 |
| 17 Characteristic of CHO cell monolayer grown in complete RPMI and CHO cell clustering in the presence of PT | 83 |
| 18 Dose response curve of CHO clustering (CC) mediated by different amounts of PT | 84 |
| 19 Results of leukocytosis-promotion (LP)-inhibition mediated by MAbPT6-2G6 | 85 |

LIST OF FIGURES (cont.)

| Figure | Page |
|--|-------------|
| 20 Representative PCR amplicons of 10B capsid protein of selected T7 phage clones that were bound by MAbPT6-2G6 | 88 |
| 21 Alignment of the M1-M6 with S1 peptide sequence | 99 |
| 22 Three dimension model of S1 subunit of pertussis toxin showing the locations of the mimotopes/ proposed epitopes recognized by MAbPT6-2G6 | 101 |
| 23 Agarose gel electrophoretic pattern of RNA extracted form MAbPT6-2G6 hybridoma cells | 103 |
| 24 Amplicons of PT6-2G6-VL and VH coding DNAs which were PCR-amplified by using <i>VL</i> and <i>VH</i> primer sets | 104 |
| 25 Agarose gel electrophoretic patterns of DNA prepared from recombinant pGEMT inserted with PT6-2G6-VL and VH coding DNA sequences | 106 |
| 26 DNA and deduced amino acid sequences of murine PT6-2G6-VL | 107 |
| 27 DNA and deduced amino acid sequences of murine PT6-2G6-VH | 108 |
| 28 Result of NCBI BLAST between amino acid sequence of VL of PT6-2G6 with murine VL in NCBI database | 109 |
| 29 Result of NCBI BLAST between amino acid sequence of VH of PT6-2G6 with murine VH in NCBI database | 110 |
| 30 Local alignment of murine PT6-2G6-VL and VH amino acid sequences to the most matched-human VL (A) and VH (B) sequences | 114 |
| 31 CDR sequences and frameworks of PT6-2G6-VL, the most matched-human VL and the expected humanized-VL (HuVL) sequences | 115 |
| 32 CDR sequences and frameworks of PT6-2G6-VH, the most matched-human VH and the expected humanized-VH (HuVH) sequences | 116 |
| 33 Amplicons used in the humanized- <i>VL</i> construction | 118 |
| 34 Amplicons of humanized-PT6-2G6-VL coding DNA | 119 |
| 35 Amplicons used in the humanized- <i>VH</i> construction | 120 |
| 36 Amplicons of humanized-PT6-2G6-VH coding DNA | 121 |

LIST OF FIGURES (cont.)

| Figure | Page |
|--|-------------|
| 37 Amplicons of PT6-2G6-HuScFv coding DNA | 123 |
| 38 The bacterial plate of transformed TG1 <i>E. coli</i> with pCANTAB5E carrying <i>PT6-2G6-huscFv</i> DNA insert | 125 |
| 39 Agarose gel electrophoretic patterns of plasmid DNA prepared from recombinant pCANTAB 5E with <i>PT6-2G-huscFv</i> DNA insert | 126 |
| 40 PCR products of pCANTAB 5E containing <i>PT6-2G-huscFv</i> DNA in TG1 <i>E. coli</i> | 127 |
| 41 Phage ELISA OD 405 of phage pools of the four sub-libraries to PT (100 ng per well) compared with BSA as negative control | 129 |
| 42 Phage ELISA of individual phage clones displaying PT6-2G6-HuScFv to 100 ng of PT compared with BSA as negative control | 130 |
| 43 Dot blot analysis of soluble proteins of HB2151 <i>E. coli</i> transfected with selected phages carrying specific PT6-2G6-HuScFv to PT by using anti-E tag monoclonal antibody and PBST as negative control | 132 |
| 44 SDS-PAGE and Western blot analysis of soluble proteins of HB2151 <i>E. coli</i> no. 1.1 and 1.3 | 133 |
| 45 SDS-PAGE and Western blot analysis of bacterial extracts of HB2151 <i>E. coli</i> PT6-2G6-HuScFv no. 1.1 and 1.3 | 135 |
| 46 Soluble HuScFv ELISA of PT6-2G6-HuScFv 1.1 and 1.3 against 0.1 µg PT compared with unrelated muScFv and diluent as ScFv negative control and blank, respectively | 137 |
| 47 DNA and deduced amino acid sequences of PT6-2G6-HuScFv | 140 |
| 48 Result of NCBI BLAST between amino acid sequence of PT6-2G6-HuVL with human VL in NCBI database | 141 |
| 49 Result of NCBI BLAST between amino acid sequences of PT6-2G6-HuVH with human VH in NCBI database | 142 |

LIST OF FIGURES (cont.)

| Figure | Page |
|---|-------------|
| 50 SDS-PAGE and Western blot analysis of PT6-2G6-HuScFv purified by HiTrap [®] Anti-E Tag column | 144 |
| 51 Western blot analysis of MAbPT6-2G6 and PT6-2G6-HuScFv to SDS PAGE-seperated PT | 146 |
| 52 Hemagglutination inhibition assay for determining the HI activity of PT6-2G6-HuScFv. | 148 |



THAMMASAT UNIVERSITY

สำนักหอสมุด