

## LIST OF TABLES

Table	Page
<b>LITERATURE REVIEW</b>	
1 Association of <i>Vibrio</i> species with different clinical syndromes	5
2 Biochemical and other characteristics of 12 <i>Vibrio</i> species found in human clinical specimens	9
3 PCR for detection of CT gene	21
<b>MATERIALS AND METHODS</b>	
4 Primers for amplification of <i>Vibrio cholerae</i> genes	52
<b>RESULTS</b>	
5 Enteric bacteria isolated from rectal swab sample of 1,066 patients with watery diarrhea who were admitted at Bamrasnaradura Infectious Disease Institute, Nonthaburi, Thailand, during April to August 2003 and February to March 2006	59
6 Comparison of the results between detection <i>V. cholerae</i> O1 in 1,066 rectal swab samples of watery diarrhea patients by MAb-based dot-ELISA and the culture method	63
7 Comparison of the results between detection <i>V. cholerae</i> O139 in 1,066 rectal swab samples of watery diarrhea patients by MAb-based dot-ELISA and the culture method	67
8 Comparison of the results between DNA amplification of <i>ctxA</i> , <i>ctxB</i> , or <i>tcpA</i> genes and standard culture method for detection of <i>V. cholerae</i> O1 and O139 in rectal swab samples	72
9 Comparison of the results between DNA amplification of <i>toxR</i> gene and standard culture method for detection of <i>V. cholerae</i> O1 and O139 in rectal swab samples	75

Table	Page
<b>APPENDIX E</b>	
10 The result of culture isolation of enteric bacterial pathogens, MAb-based dot-ELISA, and PCR from the 1,066 rectal swab samples of patients with watery diarrhea admitted at Bamrasnaradura Infectious Disease Institute, Nonthaburi, Thailand, during April to August, 2003 and February to March, 2006	103