

LIST OF TABLES

Table	Page
LITERATURES REVIEW	
1 Evaluation trials of whole <i>Leptospira</i> -based serologic tests	14
MATERIALS AND METHODS	
2 List of <i>Leptospira</i> spp. which their whole cell homogenates were used for protein extraction for 2DE	30
3 The assignment for three-color analysis-comparison of protein abundance of nine different serovars of <i>Leptospira</i> spp.	33
4 Emission filters and laser combinations	37
5 Guidelines for running 7-cm Immobiline DryStrip™ gels on Ettan IPGphor II Isoelectric Focusing Unit	57
RESULTS	
6 The yields of <i>Leptospira</i> spp. whole cell homogenates for 2D-DIGE after treatment with the 2D-Clean-up kit and protein determined using the 2D-Quant kit	73
7 2D-DIGE assignments outlined for protein abundance analysis and comparison among nine different serovars of <i>Leptospira</i> spp.	74
8 Average ratios and t-test <i>p</i> values of 11 protein spots matched in 2 groups of pathogenic <i>L. interrogans</i> (5 serovars), <i>L. borgpetersenii</i> (2 serovars) and nonpathogenic <i>L. biflexa</i> (2 serovars)	103
9 Protein identifications of spots no. 1 to 11 of <i>L. interrogans</i> serogroup Pomona serovar Pomona separated by 2DE at linear pH 3-10 (Figure 32) using MASCOT (MatrixScience)	109
10 Orthologous proteins in the database in which the peptide sequences matched with those generated from tryptic digestion and LC/MS/MS analysis of proteins in spots no. 1 to 11 of <i>L. interrogans</i> serogroup Pomona serovar Pomona separated by 2DE at linear pH 3-10	116

LIST OF TABLES (Cont.)

Table	Page
11 Orthologous proteins of the database matched with the peptide sequences generated from tryptic digestion of proteins in spots no. 1 to 22 of <i>L. borgpetersenii</i> serogroup Tarassovi serovar Tarassovi homogenate separated by 2DE at non-linear pH 3-10	134
12 Orthologous proteins of the database matched with the peptide sequences generated from tryptic digestion of proteins in spots no. 1 to 13 of homogenate of <i>L. interrogans</i> , serogroup Icterohaemorrhagiae, serovar Copenhageni separated by 2DE at non-linear pH 3-10 (Figure 46A).	137
13 Orthologous proteins of the database matched with the peptide sequences generated from tryptic digestion of proteins in spots no. 1 to 19 of homogenate of <i>L. borgpetersenii</i> , serogroup Tarassovi, serovar Tarassovi separated by 2DE at non-linear pH 3-10.	139
14 Functions of the immunogenic proteins of the two pathogenic <i>Leptospira</i> spp.	141
15 Estimated molecular masses and pI of <i>in vivo</i> expressed antigens after 2DE-immunoblotting of protein homogenate of <i>L. borgpetersenii</i> , serogroup Tarassovi, serovar Tarassovi, 30 µg/gel by using Immobiline™ DryStrip non linear pH 3-5.6 and pH 5.3–6.5 reacted to leptospirosis patients (Figures 52 -57) compared with serum from melioidosis and dengue hemorrhagic fever (DHF) (Figures 60-61)	160
16 Orthologous proteins in database matched with the peptide sequences generated from tryptic digestion and LC/MS/MS of proteins in spot no. 1 to 33 of <i>L. borgpetersenii</i> serogroup Tarassovi serovar Tarassovi separated by 2DE at non-linear pH 3-5.6 (spots no. 1-6), pH 5.3-6.5 (spots no. 7-18), pH 6.2-7.5 (spot no. 24) and non-linear pH 7-11 (spots no. 19-33) in Figure 62	163

LIST OF TABLES (Cont.)

Table		Page
17	Protein descriptions and functions of orthologous protein in Table 13	168
18	Summary of orthologous proteins	175

