

## LIST OF FIGURES

<b>Figure</b>	<b>Page</b>
<b>MATERIALS AND METHODS</b>	
1 DeCyder Differential Analysis Software co-detection and matching using the internal standard	38
2 Work chart for isotyping and sub-isotyping of ten MAb samples	53
3 Map of pcDNA3.1[+]	63
<b>RESULTS</b>	
4 The fluorescent dye labeled proteins of <i>Leptospira</i> spp.	75
5 The fluorescent dye labeled proteins of <i>L. biflexa</i> serogroup Andamana serovar Andamana and <i>L. interrogans</i> , serogroup Pomona, serovars Pomona	76
6 Statistical analyses for the comparison of protein abundance in spots by DeCyder Biological Variation Module™	77
7 DeCyder software output of spot no. 1 showing the master gel images of Cy3 and Cy5 labeled proteins	78
8 DeCyder software output of spot no. 1: three-dimensional (3D) views of the primary and secondary images localized on spot no. 1	79
9 DeCyder software output of spot no. 2 showing the master gel images of Cy3 and Cy5 labeled proteins	80
10 DeCyder software output of spot no. 2: 3D views of the primary and secondary images localized on spot no. 2	81
11 DeCyder software output of spot no. 3 showing the master gel images of Cy3 and Cy5 labeled proteins	82
12 DeCyder software output of spot no. 3: 3D views of the primary and secondary images localized on spot no. 3	83
13 DeCyder software output of spot no. 4 showing the master gel images of Cy3 and Cy5 labeled proteins	84
14 DeCyder software output of spot no. 4: 3D views of the primary and secondary images localized on spot no. 4	85

**LIST OF FIGURES (Cont.)**

<b>Figure</b>	<b>Page</b>
15 DeCyder software output of spot no. 5 showing the master gel images of Cy3 and Cy5 labeled proteins	86
16 DeCyder software output of spot no. 5: 3D views of the primary and secondary images localized on spot no. 5	87
17 DeCyder software output of spot no. 6: showing the master gel images of Cy3 and Cy5 labeled proteins	88
18 DeCyder software output of spot no. 6: 3D views of the primary and secondary images localized on spot no. 6	89
19 DeCyder software output of spot no. 7 showing the master gel images of Cy3 and Cy5 labeled proteins	90
20 DeCyder software output of spot no. 7: 3D views of the primary and secondary images localized on spot no. 7	91
21 DeCyder software output of spot no. 8 showing the master gel images of Cy3 and Cy5 labeled proteins	92
22 DeCyder software output of spot no. 8: Three-dimensional (3D) views of the primary and secondary images localized on spot no. 8	93
23 DeCyder software output of spot no. 9 showing the master gel images of Cy3 and Cy5 labeled proteins	94
24 DeCyder software output of spot no. 9: Three-dimensional (3D) views of the primary and secondary images localized on spot no. 9	95
25 DeCyder software output of spot no. 10 showing the master gel images of Cy3 and Cy5 labeled proteins	96
26 DeCyder software output of spot no. 10: Three-dimensional (3D) views of the primary and secondary images localized on spot no. 10	97
27 DeCyder software output of spot no. 11 showing the master gel images of Cy3 and Cy5 labeled proteins	98
28 DeCyder software output of spot no. 11: Three-dimensional (3D) views of the primary and secondary images localized on spot no. 11	99

## LIST OF FIGURES (Cont.)

<b>Figure</b>	<b>Page</b>
29 Graphs of logarithmic standardized protein abundance obtained from spots no. 1, 2, 3 and 4 of 2D-DIGE	100
30 Graphs of logarithmic standardized protein abundance obtained from spots no. 5, 6, 7 and 8 of 2D-DIGE	101
31 Graphs of logarithmic standardized protein abundance obtained from spots no. 9, 10 and 11 of 2D-DIGE	102
32 Proteomes of <i>Leptospira interrogans</i> , serogroup Pomona, serovar Pomona separated on Ettan DALT™ 12.5% (26 × 20 cm, 1 mm thick) gel after focusing with 18 cm Immobiline™ DryStrip gel, pH 3–10, sample load was 500 µg protein stained by Coomassie Brilliant Blue G-250 dye	105
33 MALDI-ToF MS profile of tryptic digest of 2DE separated proteins of <i>Leptospira interrogans</i> , serogroup Pomona, serovar Pomona in the gel plug of spots no. 1, 2, 3 and 4	106
34 MALDI-ToF MS profile of tryptic digest of 2DE separated proteins of <i>Leptospira interrogans</i> , serogroup Pomona, serovar Pomona in the gel plug of spots no. 5, 6, 7 and 8	107
35 MALDI-ToF MS profile of tryptic digest of 2DE separated proteins of <i>Leptospira interrogans</i> , serogroup Pomona, serovar Pomona in the gel plug of spots no. 9, 10 and 11	108
36 SDS-PAGE-separated-whole cell homogenates of various <i>Leptospira</i> spp. stained by colloidal Coomassie Brilliant Blue staining	118
37 Antigenic specificities of MAblPF1 (A) and MAblPF2 (B) revealed by SDS-PAGE and Western blot analysis against whole cell homogenates of various <i>Leptospira</i> spp.	120

## LIST OF FIGURES (Cont.)

<b>Figure</b>	<b>Page</b>
38 2D proteomic pattern of <i>L. borgpetersenii</i> serogroup Tarassovi serovar Tarassovi and 6 µg of protein samples were rehydrated (8M Urea, 0.5% IPG buffer pH 3-10NL) and IEF in individual strip holder Immobiline™ DryStrip pH 3-10 NL, 12.5% gel, stained by Silver stain	122
39 2D proteomic pattern of <i>L. borgpetersenii</i> serogroup Tarassovi serovar Tarassovi and 6 µg of protein samples were rehydrated (2M Thiourea, 7M Urea, 0.5% IPG buffer pH 3-10NL) and IEF in individual strip holder Immobiline™ DryStrip pH 3-10 NL, 12.5% gel, stained by Silver stain	123
40 2D proteomic pattern of <i>L. borgpetersenii</i> serogroup Tarassovi serovar Tarassovi. The homogenate was lysed in cell lysis buffer (8 M urea) and 6 µg of protein sample were rehydrated in rehydration buffer (8 M urea, 0.5% IPG buffer 3-10 NL and 0.2% DTT), Immobiline™ DryStrip pH 3-10NL, 12.5% gel, stained by silver stain	124
41 2D proteomic pattern of <i>L. borgpetersenii</i> serogroup Tarassovi serovar Tarassovi. The homogenate was lysed in cell lysis buffer (2 M thiourea, 7 M urea) and 6 µg of protein sample were rehydrated in rehydration buffer (2 M thiourea, 7 M urea 0.5% IPG buffer 3-10 NL and 0.2% DTT), Immobiline™ DryStrip pH 3-10 NL, 12.5% gel, stained by silver stain	125
42 2D proteomic pattern of <i>L. borgpetersenii</i> serogroup Tarassovi serovar Tarassovi. and 6 µg of protein samples were rehydrated in rehydration tray (gel side down) 12 hr and IEF in ETTANmanifold (gel side up) Immobiline™ DryStrip pH 3-10 NL, 12.5 % gel, stained by silver stain	126

## LIST OF FIGURES (Cont.)

<b>Figure</b>	<b>Page</b>
43 Proteomes of <i>Leptospira interrogans</i> , serogroup Icterohaemorrhagiae, serovar Copenhageni (A) stained by Coomassie dye. Proteins in the homogenates reacting to MAbLPF1 and MAbLPF2 in the 2DE-immunoblot are shown in B and C, respectively	130
44 Proteomes of <i>Leptospira interrogans</i> , serogroups Icterohaemorrhagiae, serovar Icterohaemorrhagiae (A) stained by Coomassie dye. Proteins in the homogenates reacting to MAbLPF1 and MAbLPF2 in the 2DE-immunoblot are shown in B and C, respectively	131
45 Proteomes of <i>Leptospira borgpetersenii</i> , serogroup Tarassovi, serovar Tarassovi (A), stained by Coomassie dye. Proteins in the homogenates reacting to MAbLPF1 and MAbLPF2 in the 2DE-immunoblot are shown in figures B and C, respectively	132
46 Proteome and immunome of whole cell homogenate of <i>L. interrogans</i> , serogroup Icterohaemorrhagiae, serovar Copenhageni (A) and <i>L. borgpetersenii</i> , serogroup Tarassovi, serovar Tarassovi (B).	133
47 Immunomes of <i>L. interrogans</i> , serogroup Icterohaemorrhagiae, serovar Copenhageni (1A, 1B), serovar Icterohaemorrhagiae (2A, 2B) and <i>L. borgpetersenii</i> , serogroup Tarassovi, serovar Tarassovi (3A, 3B). Protein in the homogenates reacting to serum of leptospirosis patient number 1, the acute phase serum (1A, 2A and 3A) was collected at day 1 <sup>st</sup> of hospitalized and convalescence phase serum (1B, 2B, 3B) was collected at day 3 <sup>rd</sup> of hospitalized.	145

## LIST OF FIGURES (Cont.)

<b>Figure</b>	<b>Page</b>
<p>48 Immunomes of <i>L. interrogans</i>, serogroup Icterohaemorrhagiae, serovar Copenhageni (1A, 1B), serovar Icterohaemorrhagiae (2A, 2B) and <i>L. borgpetersenii</i>, serogroup Tarassovi, serovar Tarassovi (3A, 3B). Protein in the homogenates reacting to serum of leptospirosis patient number 2, the acute phase serum (1A, 2A and 3A) was collected at day 1<sup>st</sup> of hospitalized and convalescence phase serum (1B, 2B, 3B) was collected at day 14<sup>th</sup> of hospitalized.</p>	146
<p>49 Immunomes of <i>L. interrogans</i>, serogroup Icterohaemorrhagiae, serovar Copenhageni (1A, 1B), serovar Icterohaemorrhagiae (2A, 2B) and <i>L. borgpetersenii</i>, serogroup Tarassovi, serovar Tarassovi (3A, 3B). Protein in the homogenates reacting to serum of leptospirosis patient number 3, the acute phase serum (1A, 2A and 3A) was collected at day 1<sup>st</sup> of hospitalized and convalescence phase serum (1B, 2B, 3B) was collected at day 14<sup>th</sup> of hospitalized.</p>	147
<p>50 Immunomes of <i>L. interrogans</i>, serogroup Icterohaemorrhagiae, serovar Copenhageni (1A, 1B), and <i>L. borgpetersenii</i>, serogroup Tarassovi, serovar Tarassovi (2A, 2B). Protein in the homogenates reacting to serum of leptospirosis patient number 4, the acute phase serum (1A, 2A and 3A) was collected at day 1<sup>st</sup> of hospitalized and convalescence phase serum (1B, 2B, 3B) was collected at day 7<sup>th</sup> of hospitalized.</p>	148
<p>51 Immunomes of <i>L. interrogans</i>, serogroup Icterohaemorrhagiae, serovar Copenhageni (1A, 1B), serovar Icterohaemorrhagiae (2A, 2B) and <i>L. borgpetersenii</i>, serogroup Tarassovi, serovar Tarassovi (3A, 3B). Protein in the homogenates reacting to serum of Dengue hemorrhagic fever patient (1A, 2A and 3A) and serum of melioidosis patient (1B, 2B, 3B).</p>	149

## LIST OF FIGURES (Cont.)

<b>Figure</b>	<b>Page</b>
<p>52 2DE-immunomic patterns (<i>In vivo</i> expressed antigens) of whole cell homogenate of <i>L. borgpetersenii</i>, serogroup Tarassovi, serovar Tarassovi separated on Immobiline™ DryStrip non linear pH 3-5.6 (1A and 1B) and pH 5.3–6.5 (2A and 2B), 12.5% polyacrylamide gel, 30 µg protein. Protein in the homogenates reacting to serum of leptospirosis patient number 1, the acute phase serum (1A and 2A) was collected at day 1<sup>st</sup> of hospitalized and convalescence phase serum (1B and 2B) was collected at day 3<sup>rd</sup> of hospitalized.</p>	150
<p>53 2DE-immunomic patterns (<i>in vivo</i> expressed antigens) of whole cell homogenate of <i>L. borgpetersenii</i>, serogroup Tarassovi, serovar Tarassovi separated on Immobiline™ DryStrip pH 6.2-7.5 (1A and 1B) and non linear pH 7-11 (2A and 2B), 12.5% polyacrylamide gel, 30 µg protein. Protein in the homogenates reacting to serum of leptospirosis patient number 1, the acute phase serum (1A and 2A) was collected at day 1<sup>st</sup> of hospitalized and convalescence phase serum (1B and 2B) was collected at day 3<sup>rd</sup> of hospitalized.</p>	151
<p>54 2DE-immunomic patterns (<i>in vivo</i> expressed antigens) of whole cell homogenate of <i>L. borgpetersenii</i>, serogroup Tarassovi, serovar Tarassovi separated on Immobiline™ DryStrip non linear pH 3-5.6 (1A and 1B) and pH 5.3–6.5 (2A and 2B), 12.5% polyacrylamide gel, 30 µg protein. Protein in the homogenates reacting to serum of leptospirosis patient number 2, the acute phase serum (1A and 2A) was collected at day 1<sup>st</sup> of hospitalized and convalescence phase serum (1B and 2B) was collected at day 14<sup>th</sup> of hospitalized.</p>	152

## LIST OF FIGURES (Cont.)

<b>Figure</b>	<b>Page</b>
<p>55 2DE-immunomic patterns (<i>in vivo</i> expressed antigens) of whole cell homogenate of <i>L. borgpetersenii</i>, serogroup Tarassovi, serovar Tarassovi separated on Immobiline™ DryStrip pH 6.2-7.5 (1A and 1B) and non linear pH 7-11 (2A and 2B), 12.5% polyacrylamide gel, 30 µg protein. Protein in the homogenates reacting to serum of leptospirosis patient number 2, the acute phase serum (1A and 2A) was collected at day 1<sup>st</sup> of hospitalized and convalescence phase serum (1B and 2B) was collected at day 14<sup>th</sup> of hospitalized.</p>	153
<p>56 2DE-immunomic patterns (<i>in vivo</i> expressed antigens) of whole cell homogenate of <i>L. borgpetersenii</i>, serogroup Tarassovi, serovar Tarassovi separated on Immobiline™ DryStrip non linear pH 3-5.6 (1A and 1B) and pH 5.3–6.5 (2A and 2B), 12.5% polyacrylamide gel, 30 µg protein. Protein in the homogenates reacting to serum of leptospirosis patient number 3, the acute phase serum (1A and 2A) was collected at day 1<sup>st</sup> of hospitalized and convalescence phase serum (1B and 2B) was collected at day 14<sup>th</sup> of hospitalized.</p>	154
<p>57 2DE-immunomic patterns (<i>in vivo</i> expressed antigens) of whole cell homogenate of <i>L. borgpetersenii</i>, serogroup Tarassovi, serovar Tarassovi separated on Immobiline™ DryStrip pH 6.2-7.5 (1A and 1B) and non linear pH 7-11 (2A and 2B), 12.5% polyacrylamide gel, 30 µg protein. Protein in the homogenates reacting to serum of leptospirosis patient number 3, the acute phase serum (1A and 2A) was collected at day 1<sup>st</sup> of hospitalized and convalescence phase serum (1B and 2B) was collected at day 14<sup>th</sup> of hospitalized.</p>	155

## LIST OF FIGURES (Cont.)

<b>Figure</b>	<b>Page</b>
<p>58 2DE-immunomic patterns (<i>in vivo</i> expressed antigens) of whole cell homogenate of <i>L. borgpetersenii</i>, serogroup Tarassovi, serovar Tarassovi separated on Immobiline™ DryStrip non linear pH 3-5.6 (1A and 1B) and pH 5.3–6.5 (2A and 2B), 12.5% polyacrylamide gel, 30 µg protein. Protein in the homogenates reacting to serum of leptospirosis patient number 4, the acute phase serum (1A and 2A) was collected at day 1<sup>st</sup> of hospitalized and convalescence phase serum (1B and 2B) was collected at day 7<sup>th</sup> of hospitalized.</p>	156
<p>59 2DE-immunomic patterns (<i>in vivo</i> expressed antigens) of whole cell homogenate of <i>L. borgpetersenii</i>, serogroup Tarassovi, serovar Tarassovi separated on Immobiline™ DryStrip pH 6.2-7.5 (1A and 1B) and non linear pH 7-11 (2A and 2B), 12.5% polyacrylamide gel, 30 µg protein. Protein in the homogenates reacting to serum of leptospirosis patient number 4, the acute phase serum (1A and 2A) was collected at day 1<sup>st</sup> of hospitalized and convalescence phase serum (1B and 2B) was collected at day 7<sup>th</sup> of hospitalized.</p>	157
<p>60 2DE-immunomic patterns (<i>in vivo</i> expressed antigens) of whole cell homogenate of <i>L. borgpetersenii</i>, serogroup Tarassovi, serovar Tarassovi separated on Immobiline™ DryStrip non linear pH 3-5.6 (1A and 1B) and pH 5.3–6.5 (2A and 2B), 12.5% polyacrylamide gel, 30 µg protein. Protein in the homogenates reacted to serum of Dengue hemorrhagic fever patient (1A, and 2A) and serum of melioidosis patient (1B and 2B).</p>	158

## LIST OF FIGURES (Cont.)

<b>Figure</b>	<b>Page</b>
61 2DE-immunomic patterns ( <i>in vivo</i> expressed antigens) of whole cell homogenate of <i>L. borgpetersenii</i> , serogroup Tarassovi, serovar Tarassovi separated on Immobiline™ DryStrip pH 6.2-7.5 (1A and 1B) and non linear pH 7-11 (2A and 2B), 12.5% polyacrylamide gel, 30 µg protein. Protein in the homogenates reacted to serum of Dengue hemorrhagic fever patient (1A, and 2A) and serum of melioidosis patient (1B and 2B).	159
62 Proteomes of <i>Leptospira borgpetersenii</i> , serogroup Tarassovi, serovar Tarassovi (A), stained by Coomassie dye, separated on Immobiline™ DryStrip non linear pH 3-5.6 (1A), pH 5.3–6.5 (2A), pH 6.2-7.5 (1B) and non linear pH 7-11 (2B), 12.5 % polyacrylamide gel, 240 µg protein.	162
63 Gradient PCR amplicons of <i>ompL1</i>	179
64 <i>Leptospira</i> DNA amplicons of eight vaccine candidate genes	180
65 Positive <i>E. coli</i> clones with inserted genes of <i>ompL1</i> of <i>Leptospira</i>	181
66 Agarose gel electrophoresis of extracted pcDNA plasmids	182
67 Indirect ELISA titers of antibodies to homogenates of <i>L. interrogans</i> serogroup Icterohaemorrhagiae serovar Icterohaemorrhagiae in the sera of vaccinated and control hamsters at different day after the booster.	183
68 Survival rate (% survival) of vaccinated and control hamsters.	184